

Randomized Controlled Community Trial of the Efficacy of a Multicomponent Stage-Matched Intervention to Increase Sun Protection among Beachgoers¹

Martin A. Weinstock, M.D., Ph.D.,*² Joseph S. Rossi, Ph.D.,†‡ Colleen A. Redding, Ph.D.,†
and Jason E. Maddock, Ph.D.§

*Dermatoepidemiology Unit, VA Medical Center, Providence, Rhode Island 02908, Department of Dermatology, Rhode Island Hospital, Providence, Rhode Island 02903, and Brown University, Providence, Rhode Island 02912; †Cancer Prevention Research Center and ‡Department of Psychology, University of Rhode Island, Kingston, Rhode Island 02881; and §Cancer Research Center of Hawaii, University of Hawaii, Honolulu, Hawaii 96822

Background. Skin cancer is about as common as all other cancers combined and is preventable by sun protection. The most intense sun exposures often occur on the beach, so we chose this setting to test an intervention to affect sun protection behaviors.

Methods. We developed a multicomponent stage-matched intervention for beachgoers and evaluated its efficacy in a randomized trial for influencing stage of change and self-reported behavior.

Results. We randomized 2,324 persons ages 16 to 65 on the beach (83% of those approached). The intervention was effective in increasing self-reported sun protective behaviors. Effects were similar across gender and age groups.

Conclusions. The beach is a good site for recruitment and intervention to prevent skin cancer in high-risk populations. Our stage-matched intervention package was effective for increasing sun protective behaviors. © 2002 American Health Foundation and Elsevier Science (USA)

Key Words: skin cancer prevention; sun exposure; sunscreen agents; health-related behaviors; health surveys; beach.

INTRODUCTION

Skin cancer incidence now roughly equals that of all other cancers combined (approximately 1.3 million cases per year) and continues to increase [1,2]. The primary cause of skin cancer is chronic unprotected exposure to the sun's ultraviolet light, as well as intense, intermittent exposures such as those that occur

at the beach [3–5]. Most skin cancers are therefore largely preventable with appropriate environmental and behavioral changes to minimize exposure. Leading public health organizations have recommended reducing sun exposure, wearing protective clothing, and using sunscreen with a sun protection factor of 15 or higher [6–8].

Health promotion efforts designed to modify lifestyle behaviors related to primary or secondary disease prevention have met with only occasional success, although interventions based on established behavioral models have tended to show the largest and most consistent effects [9]. Our intervention was based on the transtheoretical model of health behavior change [8,10]. This model suggests that individuals attempting to change a problem behavior progress through a series of five stages of change: precontemplation (not thinking about change), contemplation (seriously considering change), preparation (deciding and planning to change), action (initiation of overt behavior change), and maintenance (sustained behavior change over time). Progression from one stage to the next involves: (a) use of distinct subsets of processes (strategies and techniques) of change; (b) assessment of the importance of the pros and cons of change (decisional balance); and (c) the ability to resist situational temptations to engage in the problem behavior and increasing situational self-efficacy to engage in more healthful behaviors.

The likelihood of successful behavior change is optimized when intervention strategies are matched to the individual's stage of change [11]. Interventions tailored to stage of change have been more successful at changing behavior than control or comparison interventions across a wide range of important public health problems, including smoking cessation, dietary fat reduc-

¹ This research was supported by Grant AR43051.

² To whom correspondence and reprint requests should be addressed at Dermatoepidemiology Unit, VA Medical Center–111D, 830 Chalkstone Avenue, Providence, RI 02908-4799. Fax: 401-457-3332. E-mail: maw@brown.edu.

tion, exercise adoption, condom adoption, and adherence to mammography screening [12–23]. We sought to extend these findings to sun protection by developing an individualized, multicomponent intervention targeted to study participants' stage of change [8,24,25]. The effectiveness of this intervention package to increase sun protection was evaluated in a randomized clinical trial of at-risk beach bathers.

METHOD

Sample

Sun bathers on beaches in or near Rhode Island ($N = 2,800$) were recruited to participate in the study during the summer of 1995 and 2,324 (83% of those approached) were enrolled. A baseline sample size of 2,400 was planned for the study based on several factors, including plausible recruitment (80%) and retention rates (67%) and expected intervention effect sizes ($d = 0.20$ – 0.25). Although these study design parameters could not be known with precision a priori, reasonable estimates were available based on several years of beach pilot work [24,25] and previous population-based studies using transtheoretical model interventions for other health behaviors [17,23]. A minimum statistical power of 0.90 was used for sample size calculations, providing some protection against the possibility of smaller than expected intervention effects or greater than expected attrition rates [26–28].

Participants from ages 16 to 65 completed a survey on the beach with a trained interviewer. The sample was primarily female (60%), white (94%), single (51%), or married (40%), with at least a high school education (88%) and with a median income of \$45,000–65,000 per year. The average age of participants was 33 years ($SD = 12$), with 35% aged 16–24 (beachgoers 16–24 years of age were oversampled), 35% aged 25–39, and 29% aged 40–65. About 25% of the sample reported being current smokers and 64% reported exercising regularly at least 20 minutes a day, three times a week. Further details on the baseline characteristics of this sample have been reported elsewhere [29].

Procedures

Seven of the largest coast salt water beaches in southern Rhode Island were selected for the study. Upon receiving signed informed consent to participate, the interviewer administered the baseline survey and then randomized the participant to intervention or control conditions using a preassigned sequence. When the participant was in the treatment group, the interviewer would conduct the intervention and conclude by verifying contact information. When the participant was in the control group, the interviewer would just verify the contact information. Interviews lasted 15–25 minutes. Follow-up assessments were conducted by




FIG. 1. Study participation.

mail or telephone interview for all study participants at 2 and 12 months after baseline, and a final assessment was conducted at 24 months. Initial follow-up attempts were conducted by mail. After about 2 weeks, mail nonrespondents were contacted by telephone. We have used mixed-mode data collection procedures for most of our previous population-based intervention studies without discernable respondent bias [17,18,23]. All study participants were eligible for a lottery prize of \$1,000 for completing each assessment. Study retention rates were 83% of baseline at 2 months ($N = 1,930$), 70% of baseline at 12 months ($N = 1,628$), and 62% of baseline at 24 months ($N = 1,449$). The average retention rate was 85% of those contacted at the previous time point. No differential dropout by treatment group was observed (see Fig. 1). Additional details on study procedures including site selection and training of interviewers have been reported elsewhere [29].

Interventions

To maximize impacts on sun protective behaviors, the multicomponent intervention package was designed to be effective with individuals in each of the

five stages of change, as specified by the transtheoretical model. Interventions were designed to raise the pros and decrease the cons of sun protection, increase self-efficacy in dealing with challenging situations, and increase the frequency of use of the processes of change. Interventions delivered on the beach included an educational pamphlet, personalized sun sensitivity assessment and feedback (written and verbal), SPF 15 sunscreen, and instant sun damage imaging photographs that reveal damage to the skin that is not visible in normal light [8,24,25,30]. The pamphlet and sunscreen were generic, and the feedback and photographs were tailored to the participant. Intervention and measurement development was based on 3 years of pilot studies conducted on Rhode Island beaches [24,25,31–33].

Follow-up interventions included two three- to four-page expert system feedback reports matched to the individual's stage of change. Each report reflected the individual's current stage of change and provided information and suggestions on reducing unprotected exposure based on transtheoretical model constructs, including decisional balance, four to six stage-relevant processes of change, situational self-efficacy, and some final stage-tailored techniques an individual can use to aid in the behavior change process [34,35]. The first report was based on the data provided by participants at baseline and was mailed to participants 2 to 3 weeks after assessment. The baseline report contained normative feedback, in which an individual's change efforts on many variables were compared to that of others who had been successful at changing. The second feedback report contained both normative feedback and ipsative feedback, i.e., information on an individual's progress since his or her first report. This second report was delivered 12 months following baseline and was based on the 12-month assessment.

Additional interventions delivered by mail included (a) *Being Sun Smart*, a manual containing stage-tailored information on sun protection, and (b) a second educational pamphlet. Both interventions were delivered 8 months following baseline during the spring prior to the second summer (12 month) assessment and expert system intervention. These materials provided a booster to the previous year's intervention and provided a reminder to begin thinking about taking precautions early as the next summer approached.

The control group received no intervention. The assessments were the same for both groups.

Measures

The primary outcome measure was the nine-item Sun Protection Behavior Scale (SPBS) [29]. Each item was a self-report of behavior on a five-point Likert scale of frequency (never, rarely, sometimes, often, and always) "when in the sun for more than about 15 min-

utes." The SPBS included components of sun avoidance, sunscreen use, and hat use and has been shown to be both reliable and valid [29,36,37].

A second outcome measure was stage of change for sun protection [8,29]. Two algorithms were used to measure stage of change. Each algorithm consisted of a short series of questions designed to assess intentions and behaviors for reducing sun exposure. The purpose of the staging algorithm is to classify respondents into one of the five stages of change: precontemplation, contemplation, preparation, action, or maintenance. The general sun protection algorithm classified subjects by stage based on a combination of their intentions and behaviors to protect themselves from the sun consistently by: (a) avoiding sun exposure, (b) covering up with clothing/hats, and (c) using SPF 15 sunscreens. The sunscreen staging algorithm classified subjects by stage based on their behaviors and intentions to protect themselves from sun exposure by consistently using SPF 15 sunscreens. The development of the staging algorithms is described in more detail elsewhere [8]. For both algorithms, the precontemplation stage included participants who were not consistently protecting themselves from the sun and were not intending to start doing so within the next 12 months. The contemplation stage included participants who were not consistently protecting themselves but were seriously thinking about starting to do so within the next 12 months. Preparation stage individuals were not currently protecting themselves but were planning to start doing so within the next 30 days. Action stage individuals were protecting themselves consistently but had been doing so for fewer than 12 months. Maintenance stage individuals had been protecting themselves for 12 months or more.

Demographic variables included age, sex, income, marital status, educational level, and race/ethnicity. Brief health histories focusing on skin cancer risk factors were also obtained including sun sensitivity [30], sun exposure and sunburn history, presence of large moles, use of tanning/sun lamps [38], and personal/familial history of skin cancer.

Primary analytic techniques included *t* tests, χ^2 tests, analysis of variance, repeated-measures analysis of variance, and analysis of covariance.

This study was approved by the appropriate institutional review boards.

RESULTS

At baseline, 45% of the sample was in the precontemplation stage for general sun protection, 3% in contemplation, 14% in preparation, 4% in action, and 34% in maintenance. For sunscreen use at baseline, 56% were in precontemplation, 3% in contemplation, 10% in preparation, 5% in action, and 26% in maintenance. Strong age and gender effects were observed for both

staging algorithms [29]. Fewer than one-third of study participants were using a sunscreen with an SPF of 15 or greater at the time of the interview. Among those who were using sunscreen, about half were using a sunscreen with an SPF of 15 or greater.

Analyses were conducted to assess whether participants who did not complete the 24-month assessment differed in their sun behaviors and attitudes from those who completed the study. Study noncompleters were earlier in the stages of change, $\chi^2(4, N = 2,313) = 28.6, P < 0.001, \eta^2 = 0.012$, younger ($P < 0.001, \eta^2 = 0.017$), and less likely to use sun protection as assessed by the SPBS, $t(2,311) = 5.12, P < 0.001, \eta^2 = 0.011$. Noncompleters also reported having used tanning booths more frequently than study completers overall ($P < 0.001, \eta^2 = 0.007$) and in the past year ($P < 0.001, \eta^2 = 0.013$), were less likely to know someone with melanoma ($P < 0.001, \eta^2 = 0.007$), were more often male ($P = 0.007, \eta^2 = 0.003$), had fewer lifetime sunburns ($P = 0.013, \eta^2 = 0.005$), and spent more time outside on weekends ($P = 0.018, \eta^2 = 0.006$) and weekdays ($P = 0.029, \eta^2 = 0.005$). Study completers and noncompleters did not differ on sun sensitivity, number of severe sunburns in the past year, family history of melanoma, having any big moles, number of big moles, and general health (all P 's > 0.05 without Bonferroni correction for multiple significance testing).

Among participants who completed the 24-month assessment ($N = 1,450$), t tests and χ^2 tests were run to assess baseline differences by treatment group. No significant differences were found for age, gender, sun sensitivity, sun protection, and stage of change for sun protection. The only significant difference was in stage of change of sunscreen ($\chi^2(4, N = 1,363) = 15.3, P < 0.01$), where control group participants were slightly more likely to be in precontemplation (56.0% vs 52.6%) and maintenance (28.9% vs 25.5%) and slightly less likely to be in action (2.9% vs 6.0%) and preparation (6.5% vs 10.3%) than treatment group participants.

Sun Protection Behavior Scale

Both the intervention and the control groups increased their sun protection behaviors $F(2, 1,287) = 116.36, P < 0.001, \eta^2 = 0.153$, but the improvement in the intervention group was double that of the control group at 24 months. The group (intervention vs control) by time interaction was significant, $F(2, 1,287) = 9.95, P < 0.001, \eta^2 = 0.015$. Figure 2 displays these changes over time. Each of the three subscales of the index (sun avoidance, sunscreen use, and hat use) also demonstrated significant increases in sun protective behavior for intervention subjects relative to controls from baseline to the 24-month follow-up assessment (see Table 1).

We confirmed the independent effects of the inter-




FIG. 2. Sun Protection Behavior Scale by group over time. Scores have been adjusted for initial baseline differences. Only participants with data at all three time points are included; analysis used repeated-measures analysis of variance.

vention on behavior at 24 months ($P < 0.001$) with separate analyses of covariance that controlled for age, gender, sun sensitivity, use of tanning booths in the past year, knowing anyone with skin cancer, and socioeconomic status. All of these variables were significant predictors of sun protection at baseline [29]. However, all of these analyses except gender also revealed significant effects of the covariates on the changes in behavior over the period of study. Further evaluation using repeated-measures analyses of covariance revealed differences in effect sizes across covariates. These suggested that the intervention was most effective for younger individuals, people who had low sun sensitivity, and individuals with incomes less than \$25,000.

Stages of Change

Among those who were in the preaction (precontemplation, contemplation, and preparation) stages at baseline, individuals in the intervention group were more likely than controls to have progressed to the action or maintenance stages for reducing unprotected sun exposure at 12 ($P = 0.049$) and 24 months ($P = 0.054$) and for sunscreen use at 12 ($P = 0.001$) and 24 months ($P = 0.001$). Results are shown in Table 2 along with the results for age groups and gender. Although all of our predictions were directional (treatment group $>$ control group), we report here two-sided P values to be conservative.

We also examined the effect of the intervention on progressive advancement to any stage of change, whether or not the action stage was reached. The mean stage of people in the intervention group increased significantly more than that in the control group (see Table 1). Individuals in the intervention group were also more likely to progress to more advanced stages of change for both general staging and sunscreen staging at both 12 and 24 months. Results are reported in Table 3 along with the results for age groups and gender.

TABLE 1
Intervention Effect on Sun Protection at 12 and 24 Months

Variable	Means (SD)			P	η^2
	Baseline	12 months	24 months		
SPBS				0.001	0.015
Intervention	2.82 (0.87)	3.04 (0.82)	3.18 (0.86)		
Control	2.78 (0.88)	2.96 (0.85)	3.02 (0.85)		
Sunscreen use				0.001	0.010
Intervention	2.98 (1.28)	3.18 (1.23)	3.36 (1.24)		
Control	2.96 (1.26)	3.07 (1.23)	3.15 (1.24)		
Sun avoidance				0.008	0.008
Intervention	2.71 (0.86)	2.94 (0.82)	3.04 (0.87)		
Control	2.72 (0.86)	2.87 (0.84)	2.92 (0.85)		
Hat				0.047	0.005
Intervention	2.13 (1.14)	2.17 (1.10)	2.28 (1.19)		
Control	2.23 (1.18)	2.24 (1.15)	2.24 (1.15)		
General stage				0.004	0.009
Intervention	2.97 (1.78)	2.93 (1.78)	3.18 (1.81)		
Control	2.96 (1.80)	2.88 (1.82)	2.87 (1.84)		
Sunscreen stage				0.001	0.011
Intervention	2.54 (1.76)	2.73 (1.78)	2.87 (1.85)		
Control	2.68 (1.81)	2.61 (1.80)	2.67 (1.85)		
Age ^a					
16–24				0.004	0.027
Intervention	2.47 (0.81)	2.85 (0.82)	2.95 (0.84)		
Control	2.46 (0.78)	2.65 (0.81)	2.71 (0.79)		
25–39				0.046	0.014
Intervention	2.82 (0.86)	2.98 (0.83)	3.13 (0.86)		
Control	2.93 (0.82)	3.06 (0.81)	3.11 (0.84)		
40–65				0.065	0.012
Intervention	3.11 (0.86)	3.24 (0.77)	3.40 (0.83)		
Control	3.06 (0.90)	3.13 (0.84)	3.22 (0.85)		
Sun sensitivity ^a					
Low				0.001	0.033
Intervention	2.51 (0.84)	2.82 (0.87)	3.00 (0.90)		
Control	2.51 (0.83)	2.66 (0.83)	2.74 (0.83)		
Medium				0.049	0.010
Intervention	2.86 (0.88)	3.05 (0.78)	3.14 (0.84)		
Control	2.89 (0.85)	3.03 (0.82)	3.05 (0.83)		
High				0.501	0.005
Intervention	3.26 (0.75)	3.40 (0.69)	3.55 (0.73)		
Control	3.17 (0.82)	3.25 (0.80)	3.37 (0.79)		
Use tanning booth last year? ^a					
No				0.001	0.015
Intervention	2.94 (0.87)	3.13 (0.79)	3.26 (0.85)		
Control	2.95 (0.84)	3.06 (0.82)	3.11 (0.83)		
Yes				0.110	0.018
Intervention	2.32 (0.77)	2.68 (0.85)	2.84 (0.85)		
Control	2.27 (0.78)	2.53 (0.83)	2.62 (0.84)		
Do you know anyone who has skin cancer? ^a					
No				0.007	0.016
Intervention	2.68 (0.87)	2.94 (0.81)	3.08 (0.86)		
Control	2.63 (0.84)	2.84 (0.83)	2.87 (0.84)		
Yes				0.005	0.016
Intervention	2.97 (0.87)	3.15 (0.82)	3.28 (0.85)		
Control	3.00 (0.86)	3.07 (0.85)	3.15 (0.85)		
Socioeconomic status (assessed as annual household income) ^a					
<\$25,000				0.026	0.042
Intervention	2.72 (0.92)	3.12 (0.85)	3.21 (0.97)		
Control	2.69 (0.83)	2.84 (0.87)	2.91 (0.83)		
\$25,000–54,999				0.393	0.004
Intervention	2.73 (0.86)	2.93 (0.82)	3.07 (0.82)		
Control	2.75 (0.86)	2.91 (0.86)	3.00 (0.88)		
\$55,000+				0.014	0.016
Intervention	2.91 (0.88)	3.10 (0.81)	3.23 (0.85)		
Control	2.91 (0.88)	3.03 (0.83)	3.08 (0.85)		
Gender ^a					
Female				0.031	0.008
Intervention	2.95 (0.88)	3.18 (0.84)	3.29 (0.85)		
Control	2.88 (0.88)	3.02 (0.87)	3.11 (0.86)		
Male				0.001	0.042
Intervention	2.59 (0.82)	2.80 (0.74)	2.97 (0.85)		
Control	2.71 (0.85)	2.85 (0.80)	2.85 (0.82)		

Note. Sample size with complete data at baseline, 12-month, and 24-month time points included in analyses, $N = 1,294$, repeated-measures ANOVA.

^a Means reported are for the Sun Protection Behavior Scale.

TABLE 2
Group Differences (Percentage in Action/Maintenance) for Stage of Change

	Stage of change for sun protection							
	12 months				24 months			
	<i>N</i>	Treatment (% in A/M)	Control (% in A/M)	<i>P</i>	<i>N</i>	Treatment (% in A/M)	Control (% in A/M)	<i>P</i>
All	973	25.9	20.5	0.049	845	31.5	25.5	0.054
Ages 16–24	317	23.3	15.8	0.095	261	32.3	23.4	0.071
Ages 25–39	410	24.5	24.8	0.951	357	26.3	25.8	0.921
Ages 40–65	246	31.1	19.3	0.035	227	38.2	27.9	0.100
Females	589	27.9	20.0	0.025	503	30.9	26.0	0.219
Males	384	22.8	21.4	0.732	342	32.4	24.9	0.124
	Stage of change for sunscreen use							
All	1092	22.3	13.5	0.001	948	27.1	17.0	0.001
Ages 16–24	332	20.5	11.7	0.029	278	20.9	15.3	0.223
Ages 25–39	450	19.9	14.7	0.147	393	26.1	18.9	0.090
Ages 40–65	310	27.1	13.6	0.004	277	33.5	16.0	0.001
Females	647	25.0	14.4	0.001	551	31.7	20.5	0.003
Males	445	18.3	12.0	0.065	397	20.7	12.2	0.023

Note. Only individuals in preaction at baseline are included. Two-tailed *P* values are reported.

Finally, we examined the effect of the intervention on relapse rates, individuals in action or maintenance at baseline who regressed to a preaction stage at follow-up. No significant differences between groups were found in relapse rates for reducing unprotected sun exposure (treatment = 34.6% vs control = 30.0% at 12 months; treatment = 20.7% vs control = 21.2% at 24 months) or for sunscreen use (treatment = 29.4% vs control = 28.8% at 12 months; treatment = 26.6% vs control = 27.3% at 24 months).

DISCUSSION

Few previous intervention studies have been effective at reducing unprotected sun exposure behaviors, particularly on a population basis. Typical intervention programs are relatively brief in duration, adopt minimal or “one-shot” intervention strategies, are not targeted to entire populations, and are based on limited or no theoretical framework [8]. Not surprisingly, such programs at best demonstrate increases in knowledge and changes in attitudes but not changes in behaviors. Some more intensive and theoretically guided intervention approaches have been more successful (e.g. [39], 1,685 [46], [40]), while others have not [41]. Our study is one of the first to show demonstrable and lasting impacts on protective sun exposure behaviors in an at-risk sample on a population basis. The magnitude of these effects was modest, as may be expected in a large-scale, population-based public health intervention in which the study sample reflects the high proportion of people in the precontemplation stage that is found in the general population. Significant results were obtained for behavioral outcomes—including self-reported frequency of sunscreen use, sun avoidance,

and wearing hats—and for stages of change. Stage of change results included both progression to the action stage and any forward stage progression and were obtained for both general sun protection behavior and a specific and salient marker behavior, sunscreen use. Increases in protective behavior at the end of treatment (12 months) were maintained and even increased at follow-up (24 months). These results support our high level of tailoring and retailoring on the variables: stages of change, decisional balance, self-efficacy, and processes of change characteristic of transtheoretical model-based interventions [34,35,42].

These results are consistent with two recent studies reporting the efficacy of sun protection expert system feedback (one component of this intervention) as part of a multiple behavioral risk intervention package in two different samples [42,43]. These results also add to the supportive evidence of previous studies demonstrating the effectiveness of transtheoretical model-based or stage-tailored interventions for a range of different health behaviors (e.g., [12–17,19–23,42–45]).

Limitations of our study include reliance on self-reported outcomes, which allow for possible bias, particularly social desirability bias, or unbiased yet inaccurate responses. Although we found an effect of our overall intervention package, we could not differentiate the effects of its various components with this research design. We also could not assess outcome in the dropouts, so a complete intention-to-treat analysis may have found smaller effects.

Several aspects of these results are noteworthy. In particular, it is especially encouraging that the intervention not only moved individuals to the action stage but also accelerated all forward movement through the

REFERENCES

1. Miller DL, Weinstock MA. Nonmelanoma skin cancer in the United States: incidence. *J Am Acad Dermatol* 1994;30:774-8.
2. Wingo PA, Ries LAG, Rosenberg HM, Miller DS, Edwards BK. Cancer incidence and mortality, 1973-1995: a report card for the U.S. *Cancer* 1998;82:1197-207.
3. Council on Scientific Affairs. Harmful effects of ultraviolet radiation. *JAMA* 1989;262:380-4.
4. DiGiovanna JJ. Prevention of skin cancer due to sun exposure. In: Devita VJ, Hellman S, Rosenberg S, editors. *Cancer prevention*. Philadelphia: Lippincott, 1991:1-9.
5. Weinstock MA, Clark JW, Calabresi P. Melanoma. In: Calabresi P, Schein PS, editors. *Medical oncology: basic principles and clinical management of cancer*, 2nd ed. New York: McGraw-Hill, 1993;545-63.
6. Koh HK, Lew RA. Sunscreens and melanoma: implications for prevention. *J Natl Cancer Inst* 1994;86:78-9.
7. Rossi JS. The hazards of sunlight: a report on the Consensus Development Conference on sunlight, ultraviolet radiation and the skin. *Health Psychol* 1989;11(3):4-6.
8. Rossi JS, Blais LM, Redding CA, Weinstock MA. Behavior change for reducing sun and ultraviolet light exposure: implications for interventions. *Dermatol Clin* 1995;13:613-22.
9. Glanz K, Lewis FM, Rimer BK, editors. *Health behavior and health education: theory, research, and practice*, 2nd ed. San Francisco: Jossey-Bass, 1997.
10. Prochaska JO, Redding CA, Evers KE. The transtheoretical model and stages of change. In: Glanz K, Lewis FM, Rimer BK, editors. *Health behavior and health education*, 2nd ed. San Francisco: Jossey-Bass, 1997:60-84.
11. Redding CA, Rossi JS, Rossi SR, Velicer WF, Prochaska JO. Health behavior models. In: Hyner GC, Peterson KW, Travis JW, Dewey JE, Foerster JJ, Frammer EM, editors. *SPM handbook of health assessment tools*. Pittsburgh: The Society of Prospective Medicine & The Institute for Health and Productivity Management, 1999:83-93.
12. Campbell MK, DeVellis BM, Strecher VJ, Ammerman AS, DeVellis RF, Sandler RS. Improving dietary behavior: the effectiveness of tailored messages in primary care settings. *Am J Public Health* 1994;84:783-7.
13. The CDC AIDS Community Demonstration Projects Research Group. Community level HIV intervention in 5 cities: final outcome data from the CDC AIDS Community Demonstration Projects. *Am J Public Health* 1999;89:336-45.
14. Marcus BH, Banspach SW, Lefebvre RC, Rossi JS, Carleton RA, Abrams DB. Using the stages of change model to increase the adoption of physical activity among community participants. *Am J Health Promot* 1992;6:424-9.
15. Marcus BH, Bock BC, Pinto BM, Forsyth LH, Roberts MB, Traficante RM. Efficacy of an individualized, motivationally-tailored physical activity intervention. *Ann Behav Med* 1998;20:174-80.
16. Prochaska JO, DiClemente CC, Velicer WF, Rossi JS. Standardized, individualized, interactive and personalized self-help programs for smoking cessation. *Health Psychol* 1993;12:399-405.
17. Prochaska JO, Velicer WF, Fava JL, Rossi JS, Tsoh JY. Evaluating a population-based recruitment approach and a stage-based expert system intervention for smoking cessation. *Addict Behav* 2001;26:583-602.
18. Prochaska JO, Velicer WF, Fava JL, *et al.* Counselor and stimulus control enhancements of a stage-matched expert system intervention for smokers in a managed care setting. *Prev Med* 2001;32:23-32.
19. Rakowski WR, Ehrich B, Goldstein MG, *et al.* Increasing mammography among women aged 40-74 by use of a stage-matched, tailored intervention. *Prev Med* 1998;27:748-56.
20. Redding CA, Prochaska JO, Rossi JS, *et al.* Effectiveness of a randomized clinical trial of stage-matched interventions targeting condom use and smoking in young urban females. *Ann Behav Med* 2002;24.
21. Schneider Jamner M, Wolitski RJ, Corby NH. Impact of a longitudinal community HIV intervention targeting injecting drug users' stages of change for condom use and bleach. *Am J Health Promot* 1997;12:15-24.
22. Skinner CS, Strecher VJ, Hospers HJ. Physician recommendations for mammography: do tailored messages make a difference? *Am J Public Health* 1994;84:43-9.
23. Velicer WF, Prochaska JO, Fava JL, Laforge RG, Rossi JS. Interactive versus non-interactive interventions and dose-response relationships for stage matched smoking cessation programs in a managed care setting. *Health Psychol* 1999;18:21-8.
24. Rossi JS, Blais LM, Weinstock MA. The Rhode Island Sun Smart Project: skin cancer prevention reaches the beaches. *Am J Public Health* 1994;84:672-4.
25. Weinstock MA, Rossi JS. The Rhode Island Sun Smart Project: a scientific approach to skin cancer prevention. *Clin Dermatol* 1998;16:411-3.
26. Cohen J. *Statistical power analysis for the behavioral sciences*, 2nd ed. Hillsdale (NJ): Erlbaum, 1988.
27. Maddock JE, Rossi JS. Statistical power of articles published in three health psychology related journals. *Health Psychol* 2001;20:76-8.
28. Rossi JS. Statistical power in psychological research: what have we gained in 20 years? *J Consult Clin Psychol* 1990;58:646-56.
29. Weinstock MA, Rossi JS, Redding CA, Maddock JE, Cottrill SD. Sun protection behaviors and stages of change for the primary prevention of skin cancers among beachgoers in southeastern New England. *Ann Behav Med* 2000;22:286-93.
30. Weinstock MA. Assessment of sun sensitivity by questionnaire: validity of items and formulation of a prediction rule. *J Clin Epidemiol* 1992;45:547-52.
31. Rossi JS, Blais LM. Readiness to reduce sun exposure to prevent skin cancer. *Int J Psychol* 1992;27:628.
32. Rossi JS, Blais LM, Weinstock MA. Skin cancer prevention at the beach. *Ann Behav Med* 1993;15(suppl):87.
33. Rossi JS, Redding CA, Weinstock MA. Technology based interventions for skin cancer prevention. *Ann Behav Med* 1998;20(suppl):S180.
34. Redding CA, Prochaska JO, Pallonen UE, *et al.* Transtheoretical individualized multimedia expert systems targeting adolescents' health behaviors. *Cog Behav Pract* 1999;6:144-53.
35. Velicer WF, Prochaska JO, Bellis JM, *et al.* An expert system intervention for smoking cessation. *Addict Behav* 1993;18:269-90.
36. Maddock JE, Rossi JS, Redding CA, Meier KS, Velicer WF, Prochaska JO. Development of transtheoretical model constructs for sun protection behaviors among adolescents. *Ann Behav Med* 1998;20(suppl):S208.
37. Rossi JS, Redding CA, Blais LM, Weinstock MA. Sun, surf, and science: the Rhode Island Sun Smart Program. *Ann Behav Med* 1995;17(suppl):S064.
38. Karagas MR, Stannard VA, Mott LA, Slattery MJ, Spencer SK, Weinstock MA. Use of tanning devices and risk of basal cell and squamous cell skin cancers. *J Natl Cancer Inst* 2002;94:224-6.

39. Dietrich AJ, Olson AL, Sox CH, Tosteson TD, Grant-Petersson J. Persistent increase in children's sun protection in a randomized controlled community trial. *Prev Med* 2000;31:569-74.
40. Lombard D, Neubauer TE, Canfield D, Winett RA. Behavioral community intervention to reduce the risk of skin cancer. *J Appl Behav Anal* 1991;24:677-86.
41. Mermelstein R, Weeks K, Turner L, Cobb J. When tailored feedback backfires: a skin cancer prevention intervention for adolescents. *Cancer Res Ther Control* 2002;8:69-79.
42. Prochaska JO, Velicer WF, Rossi JF, *et al.* Impact of simultaneous stage-matched expert systems for multiple behaviors in a population of parents. *Ann Behav Med* 2002; in press.
43. Redding CA, Prochaska JO, Goldstein M, *et al.* Efficacy of stage-matched expert systems in primary care patients to decrease smoking, dietary fat, sun exposure, and relapse from mammography. *Ann Behav Med* 2002;24(suppl):S191.
44. Glanz K, Patterson RE, Kristal AR, *et al.* Impact of work site health promotion on stages of dietary change: the Working Well trial. *Health Educ Behav* 1998;25:448-63.
45. Hollis JF, Polen M, Whitstock E, Velicer WF, Redding CA, Lichtenstein E. Efficacy of a brief tobacco prevention and cessation program for teens seen in routine medical care (TEEN REACH). *Ann Behav Med* 2002;24(suppl):S107.
46. Glanz K, Geller AC, Shigaki D, Maddock JE, Isneec MR. A randomized trial of skin cancer prevention in aquatics settings: the Pool Cool program. *Health Psych*, in press.